

Clinique dentaire Dr Jacques Vaillancourt et Dre Nadia Fournier

APICOECTOMY POST-OPERATION ADVICE

You should apply ice to the area for 10 to 12 hours after the surgery, and rest during that time.

There could be swollen to the gum, lip or cheek, this is normal for the first 48 hours. The swollen will disappear within 3-6 days.

If there is redness tint in your saliva for the first 2-3 days, don't worry it is normal.

Smoking slows down the healing process; avoid smoking as long as you can.

Your stitches will dissolve on its own after a month. We prefer to have a control visit 10-14 days after the surgery to take the stitches out.

Avoid pulling on your lip or on the area where the surgery took place in order to see your tooth.

Any discomfort or pain will completely disappear after 2 weeks.

Analgesic extra-strength (Tylenol) should sooth you after the surgery.

For the first week soft food will be your diet.

Avoid brushing the area or rinsing vigorously

RECOMMENDATIONS :

---

---

---

Dr Jacques Vaillancourt

Dre Nadia Fournier

[www.dentistevaillancourt.com](http://www.dentistevaillancourt.com)



[dentiste@dentistevaillancourt.com](mailto:dentiste@dentistevaillancourt.com)



dentiste vaillancourt et fournier

73 rue de l'Église, Windsor, Qc, J1S 2A5



819-845-3080